

Biorthogonal Wavelet Packets Associated with Nonuniform Multiresolution

Firdous Ahmad Shah

¹ Department of Mathematics, University of Kashmir, South Campus, Anantnag-192101, Jammu and Kashmir, India. E-mail: fashah79@gmail.com
(Received February 2, 2013, accepted October 13, 2013)

Abstract. In this paper, we introduce the notion of biorthogonal wavelet packets associated with nonuniform multiresolution analysis and study their characteristics by means of Fourier transform. Three biorthogonal formulas regarding these wavelet packets are established. Moreover, it is shown how to obtain several new Riesz bases of the space $L^2(\mathbb{R})$ by constructing a series of subspaces of these nonuniform wavelet packets.

Keywords: Wavelet; nonuniform multiresolution analysis; biorthogonal wavelet packet; Riesz basis; Fourier transform.

1. Introduction

In his pioneering paper, Mallat [11] first formulated a new and remarkable idea of multiresolution analysis (MRA) which deals with a general formalism for the construction of an orthonormal basis of wavelet bases. Mathematically, an MRA consists of a sequence of embedded closed subspaces, $\{V_j : j \in \mathbb{Z}\}$ of $L^2\mathbb{R}$ such that $f \in V_j$ if and only if $f \in 2V_{j+1}$. Furthermore, there exists an element $\varphi \in V_0$ such that the collection of integer translates of function φ , $\{\varphi(x - k) : k \in \mathbb{Z}\}$ represents a complete orthonormal system for V_0 . The function φ is called the *scaling function* or the *father wavelet*. Recently, the idea of MRA and wavelets have been generalized in many different settings, for example, one can replace the dilation factor 2 by an integer $M \geq 2$ and in higher dimensions, it can be replaced by a dilation matrix A , in which case the number of wavelets required is $|\det A| - 1$. But in all these cases, the translation set is always a group. In the two papers [6, 7], Gabardo and Nashed considered a generalization of Mallat's [11] celebrated theory of MRA based on spectral pairs, in which the translation set acting on the scaling function associated with the MRA to generate the subspace V_0 is no longer a group, but is the union of \mathbb{Z} and a translate of \mathbb{Z} . More precisely, this set is of the form $\Lambda = \{0, r/N\} + 2\mathbb{Z}$, where $N \geq 1$ is an integer, $1 \leq r \leq 2N - 1$, r is an odd integer relatively prime to N . They call this a *nonuniform multiresolution analysis* (NUMRA). Moreover, they have provided the necessary and sufficient conditions for the existence of associated wavelets in $L^2(\mathbb{R})$. Later on, Gabardo and Yu [8, 9] continued their research in this non-standard setting and gave the characterization of all nonuniform wavelets associated with nonuniform multiresolution analysis.

It is well-known that the classical orthonormal wavelet bases have poor frequency localization. To overcome this disadvantage, Coifman *et al.* [5] constructed univariate orthogonal wavelet packets as a generalization of Walsh systems. Wavelet packets give rise to a large class of orthonormal bases of $L^2(\mathbb{R})$, each one corresponding to a different splitting of $L^2(\mathbb{R})$ into direct sum of its closed subspaces. Wavelet packets, due to their nice characteristics have been widely applied to signal processing, coding theory, image compression, fractal theory and solving integral equations and so on. Chui and Li [3] generalized the concept of orthogonal wavelet packets to the case of non-orthogonal wavelet packets so that they can be applied to the spline wavelets and so on. The introduction of biorthogonal wavelet packets attributes to Cohen and Daubechies [4]. Shen [14] generalized the notion of univariate orthogonal wavelet packets to the case of multivariate wavelet packets. Other notable generalizations are the orthogonal version of vector-valued wavelet packets [2], higher dimensional wavelet packets with arbitrary dilation matrix [10], the orthogonal p -wavelet packets and the p -wavelet frame packets on the positive half-line \mathbb{R}^+ [12, 13].

In his recent paper, Behera [1] has constructed nonuniform wavelet packets associated with nonuniform multiresolution analysis. He proved lemmas on the so-called splitting trick and several theorems

concerning the construction of nonuniform wavelet packets on \mathbb{R} . It is well known that the orthogonal wavelet packets have many desired properties such as compact support, good frequency localization and vanishing moments. However, there is no continuous symmetry which is a much desired property in image and signal processing. To achieve symmetry, several generalizations of scalar orthogonal wavelet packets have been investigated in literature. The biorthogonal wavelet packets achieve symmetry where the orthogonality is replaced by the biorthogonality. Therefore, the main goal of this paper is to introduce the notion of biorthogonal wavelet packets associated with nonuniform multiresolution analysis and investigate their properties by means of the Fourier transform. Further, we also construct several new Riesz bases of space $L^2(\mathbb{R})$ by constructing a series of subspaces of nonuniform wavelet packets.

2. Nonuniform multiresolution analysis and the wavelet packets

In this section, we state some basic preliminaries and definitions including nonuniform multiresolution analysis, the associated nonuniform wavelets and wavelet packets. More details are referring to [6–9].

Definition 2.1. We say that a pair of functions $f(x), \tilde{f}(x) \in L^2(\mathbb{R})$ are *biorthogonal*, if their translates satisfy

$$\langle f(\cdot), \tilde{f}(\cdot - \lambda) \rangle = \delta_{0,\lambda}, \quad \lambda \in \Lambda, \quad (2.1)$$

where $\delta_{0,\lambda}$ is *Kronecker symbol*, i.e., $\delta_{0,\lambda} = 1$ when $\lambda = 0$ and $\delta_{0,\lambda} = 0$, otherwise.

Definition 2.2. Let \mathbb{H} be a Hilbert space. A sequence $\{f_k\}_{k=1}^{\infty}$ of \mathbb{H} is said to be a Riesz basis for \mathbb{H} if there exist constants A and $B, 0 < A \leq B < \infty$ such that any $f \in \mathbb{H}$ can be represented as a series $f = \sum_{k=1}^{\infty} c_k f_k = 1$ converging in \mathbb{H} with

$$A \|f\|_2^2 \leq \sum_{k=1}^{\infty} |c_k|^2 \leq B \|f\|_2^2. \quad (2.2)$$

We first recall the definition of nonuniform multiresolution analysis and some of its properties.

Definition 2.3. A sequence $\{V_j : j \in \mathbb{Z}\}$ of closed subspaces $L^2(\mathbb{R})$ is called a *nonuniform multiresolution analysis* of $L^2(\mathbb{R})$ if the following hold:

- (i) $V_j \subset V_{j+1}$, for all $j \in \mathbb{Z}$,
- (ii) $\cup_{j \in \mathbb{Z}} V_j$ is dense in $L^2(\mathbb{R})$ and $\cap_{j \in \mathbb{Z}} V_j = \{0\}$,
- (iii) $f(x) \in V_j$ if and only if $f(2N \cdot) \in V_{j+1}$,
- (iv) there is a function φ in V_0 , called the *scaling function*, such that the system of functions $\{\varphi(\cdot - \lambda) : \lambda \in \Lambda\}$ forms a Riesz basis for subspace V_0 .

It is worth noticing that, when $N = 1$, one recovers from the definition above the standard definition of a one-dimensional multiresolution analysis with dilation factor equal to 2. When $N > 1$, the dilation factor of $2N$ ensures that $2N\Lambda \subset 2\mathbb{Z} \subset \Lambda$.

Since $\varphi(x) \in V_0 \subset V_1$, there exists sequence $\{a_\lambda\}_{\lambda \in \Lambda}$ with $\sum_{\lambda \in \Lambda} |a_\lambda|^2 < \infty$ such that

$$\varphi\left(\frac{x}{2N}\right) = \sum_{\lambda \in \Lambda} a_\lambda \varphi(x - \lambda). \quad (2.3)$$

Taking Fourier transform of (2.3), we get

$$\hat{\varphi}(2N\xi) = h_0(\xi) \hat{\varphi}(\xi), \tag{2.4}$$

where $h_0(\xi) = \sum_{\lambda \in \Lambda} a_\lambda e^{-2\pi i \lambda \xi}$, is called the *symbol* of $\varphi(x)$.

Let $W_j : j \in \mathbb{Z}$ be the direct complementary subspaces of V_j in V_{j+1} . Assume that there exist a set of $2N - 1$ functions $\{\psi_1, \psi_2, \dots, \psi_{2N-1}\}$ in $L^2(\mathbb{R})$ such that their translates and dilations form a Riesz basis of W_j , i.e,

$$W_j = \overline{\text{span}} \{ \psi_r((2N)^j \cdot -\lambda) : \lambda \in \Lambda, 1 = 1, 2, \dots, 2N - 1 \}, j \in \mathbb{Z}. \tag{2.5}$$

Since $\psi_r(x) \in W_0 \subset V_1, 1 \leq r \leq 2N - 1$, there exists a sequence $\{a_\lambda^r\}_{\lambda \in \Lambda}$ with $\sum_{\lambda \in \Lambda} |a_\lambda^r|^2 < \infty$ such that

$$\frac{1}{2N} \psi_r\left(\frac{x}{2N}\right) = \sum_{\lambda \in \Lambda} a_\lambda^r \varphi(x - \lambda). \tag{2.6}$$

Implementing the Fourier transform for both sides of (2.6) gives

$$\hat{\psi}_r(2N\xi) = h_r(\xi) \hat{\varphi}(\xi), \tag{2.7}$$

where

$$h_r(\xi) = \sum_{\lambda \in \Lambda} a_\lambda^r e^{-2\pi i \lambda \xi}, 1 \leq r \leq 2N - 1. \tag{2.8}$$

If $\varphi(x), \tilde{\varphi}(x) \in L^2(\mathbb{R})$ are a pair of *biorthogonal scaling functions*, then it follows by Definition 2.1 that

$$\langle \varphi(\cdot), \tilde{\varphi}(\cdot - \lambda) \rangle = \delta_{0,\lambda}, \lambda \in \Lambda. \tag{2.9}$$

Moreover, we say that $\psi_r(x), \tilde{\psi}_r(x) \in L^2(\mathbb{R}), 1 \leq r \leq 2N - 1$ are pair of biorthogonal nonuniform wavelets associated with a pair of biorthogonal scaling functions $\varphi(x)$ and $\tilde{\varphi}(x)$ if the family $\{\psi_r(\cdot - \lambda) : \lambda \in \Lambda, 1 = 1, \dots, 2N - 1\}$ is a Riesz basis of W_0 , and

$$\langle \varphi(\cdot), \tilde{\psi}_r(\cdot - \lambda) \rangle = 0, \lambda \in \Lambda, 1 \leq r \leq 2N - 1, \tag{2.10}$$

$$\langle \tilde{\varphi}(\cdot), \psi_r(\cdot - \lambda) \rangle = 0, \lambda \in \Lambda, 1 \leq r \leq 2N - 1, \tag{2.11}$$

$$\langle \psi_r(\cdot), \tilde{\psi}_s(\cdot - \lambda) \rangle = \delta_{r,s} \delta_{0,\lambda}, \lambda \in \Lambda, 1 \leq r, s \leq 2N - 1. \tag{2.12}$$

Set

$$W_j^r = \overline{\text{span}} \{ \psi_r((2N)^j \cdot -\lambda) : \lambda \in \Lambda \}, j \in \mathbb{Z}, 1 \leq r \leq 2N - 1. \tag{2.13}$$

By definition of W_j and formulae (2.9)–(2.12), we obtain the following proposition.

Proposition 2.4. If $\psi_r(x), \tilde{\psi}_r(x) \in L^2(\mathbb{R}), 1 \leq r \leq 2N - 1$, are a pair of biorthogonal nonuniform wavelets associated with a pair of biorthogonal scaling functions $\varphi(x), \tilde{\varphi}(x)$, then

$$L^2(\mathbb{R}) = \bigoplus_{j \in \mathbb{Z}} W_j = \bigoplus_{j \in \mathbb{Z}} \bigoplus_{r=1}^{2N-1} W_j^r, \tag{2.14}$$

where \bigoplus denotes the direct sum.

Similar to (2.3) and (2.6), there exist finite sequences $\{\tilde{a}_\lambda\}_{\lambda \in \Lambda}$ and $\{\tilde{a}_\lambda^r\}_{\lambda \in \Lambda}$, $1 \leq r \leq 2N - 1$ such that $\tilde{\varphi}(x)$ and $\tilde{\psi}_r$ satisfy the following equations:

$$\tilde{\varphi}\left(\frac{x}{2N}\right) = \sum_{\lambda \in \Lambda} \tilde{a}_\lambda \tilde{\varphi}(x - \lambda). \quad (2.15)$$

$$\frac{1}{2N} \tilde{\psi}_r\left(\frac{x}{2N}\right) = \sum_{\lambda \in \Lambda} \tilde{a}_\lambda^r \tilde{\varphi}(x - \lambda). \quad (2.16)$$

For $n = 0, 1, 2, \dots$, the basic *biorthogonal nonuniform wavelet packets* ω_n and $\tilde{\omega}_n$ (as defined in [1]) associated with the scaling functions $\varphi(x)$ and $\tilde{\varphi}(x)$, respectively, are defined recursively by

$$\omega_{q+2Np}(x) = \sum_{\lambda \in \Lambda} (2N) a_\lambda^q \omega_p(2Nx - \lambda), \quad (2.17)$$

$$\tilde{\omega}_{q+2Np}(x) = \sum_{\lambda \in \Lambda} (2N) \tilde{a}_\lambda^q \tilde{\omega}_p(2Nx - \lambda), \quad (2.18)$$

where $p \geq 0$ is the unique element such that $n = q + 2Np$, $0 \leq q \leq 2N - 1$ holds.

Applying the Fourier transform for the both sides of (2.17) and (2.18) yields, respectively,

$$\hat{\omega}_{q+2Np}(x) = h_q((2N)^{-1}\xi) \hat{\omega}_p((2N)^{-1}\xi), \quad 0 \leq q \leq 2N - 1 \quad (2.19)$$

$$\hat{\tilde{\omega}}_{q+2Np}(x) = \tilde{h}_q((2N)^{-1}\xi) \hat{\tilde{\omega}}_p((2N)^{-1}\xi), \quad 0 \leq q \leq 2N - 1. \quad (2.20)$$

Lemma 2.5[6]. Let $\varphi(x), \tilde{\varphi}(x)$ be a pair of scaling functions. Then $\varphi(x), \tilde{\varphi}(x)$ are biorthogonal if and only if

$$\sum_{\lambda \in \Lambda} \hat{\varphi}(\xi - \lambda) \overline{\hat{\tilde{\varphi}}(\xi - \lambda)} = 1, \quad a. e. \xi \in \mathbb{R}.$$

3. Properties of biorthogonal wavelet packets

In this section, we characterize the biorthogonality property of the nonuniform wavelet packets by means of Fourier transform.

Lemma 3.1. Assume that $\omega_q, \tilde{\omega}_q \in L^2(\mathbb{R})$, $1 \leq q \leq 2N - 1$ are a pair of biorthogonal nonuniform wavelets associated with a pair of biorthogonal scaling functions ω_0 and $\tilde{\omega}_0$. Then, we have

$$\sum_{\sigma=0}^{2N-1} h_p((2N)^{-1}(\xi + 2\pi\sigma)) \overline{\tilde{h}_q((2N)^{-1}(\xi + 2\pi\sigma))} = \delta_{p,q}, \quad 0 \leq p, q \leq 2N - 1. \quad (3.1)$$

Proof. By using equations (2.9)–(2.13), (2.19), (2.20) and Lemma 2.5, we obtain

$$\begin{aligned}
 \delta_{p,q} &= \sum_{\lambda \in \Lambda} \omega_p(\xi + 2\pi\lambda) \overline{\tilde{\omega}_q(\xi + 2\pi\lambda)} \\
 &= \sum_{\lambda \in \Lambda} h_p((2N)^{-1}(\xi + 2\pi\lambda)) \widehat{\omega}_0((2N)^{-1}(\xi + 2\pi\lambda)) \\
 &\quad \times \overline{\widehat{\tilde{\omega}}_0((2N)^{-1}(\xi + 2\pi\lambda))} \overline{\tilde{h}_q((2N)^{-1}(\xi + 2\pi\lambda))} \\
 &= \sum_{\sigma=0}^{2N-1} h_p((2N)^{-1}(\xi + 2\pi\sigma)) \overline{\tilde{h}_q((2N)^{-1}(\xi + 2\pi\sigma))} \\
 &\quad \times \sum_{\lambda \in \Lambda} \widehat{\omega}_0((2N)^{-1}(\xi + 2\pi\sigma) + 2\pi\lambda) \overline{\widehat{\tilde{\omega}}_0((2N)^{-1}(\xi + 2\pi\sigma) + 2\pi\lambda)} \\
 &= \sum_{\sigma=0}^{2N-1} h_p((2N)^{-1}(\xi + 2\pi\sigma)) \overline{\tilde{h}_q((2N)^{-1}(\xi + 2\pi\sigma))}. \quad \square
 \end{aligned}$$

Theorem 3.2. Suppose $\{\omega_n(x) : n \geq 0\}$ and $\{\tilde{\omega}_n(x) : n \geq 0\}$ are nonuniform wavelet packets with respect to a pair of biorthogonal scaling functions $\omega_0(x)$ and $\tilde{\omega}_0(x)$, respectively. Then, for $n \geq 0$, we have

$$\langle \omega_n(\cdot), \tilde{\omega}_n(\cdot - \lambda) \rangle = \delta_{0,\lambda}, \quad \lambda \in \Lambda. \tag{3.2}$$

Proof. We prove this result by using induction on n . It follows from (2.9) and (2.12) that the claim is true for $n = 0$ and $n = 1, 2, \dots, 2N - 1$. Assume that (3.2) holds when $n < \ell$, where $\ell > 0$. Then, we prove the result (3.2) for $n = \ell$. Order $n = q + 2Np$, where $p \geq 0$, $0 \leq q \leq 2N - 1$ and $p < n$. Therefore, by induction assumption, we have

$$\langle \omega_p(\cdot), \tilde{\omega}_p(\cdot - \lambda) \rangle = \delta_{0,\lambda} \iff \sum_{\lambda \in \Lambda} \widehat{\omega}_p(\xi - \lambda) \overline{\widehat{\tilde{\omega}}_p(\xi - \lambda)} = 1, \quad \xi \in \mathbb{R}.$$

Using Proposition 2.4, Lemma 2.5, (2.19) and (2.20), we obtain

$$\begin{aligned}
 \langle \omega_n(\cdot), \tilde{\omega}_n(\cdot - \lambda) \rangle &= \langle \widehat{\omega}_n(\cdot), \widehat{\tilde{\omega}}_n(\cdot - \lambda) \rangle \\
 &= \int_{\mathbb{R}} \widehat{\omega}_{q+2Np}(\xi) \overline{\widehat{\tilde{\omega}}_{q+2Np}(\xi)} e^{2\pi i \lambda \xi} d\xi \\
 &= \int_{\mathbb{R}} h_q((2N)^{-1}\xi) \widehat{\omega}_p((2N)^{-1}\xi) \overline{\tilde{h}_q((2N)^{-1}\xi)} \overline{\widehat{\tilde{\omega}}_p((2N)^{-1}\xi)} e^{2\pi i \lambda \xi} d\xi \\
 &= \sum_{\lambda \in \Lambda} \int_{2N([0,2\pi]+\lambda)} h_q((2N)^{-1}\xi) \widehat{\omega}_p((2N)^{-1}\xi) \overline{\tilde{h}_q((2N)^{-1}\xi)} \overline{\widehat{\tilde{\omega}}_p((2N)^{-1}\xi)} e^{2\pi i \lambda \xi} d\xi \\
 &= \int_{2N[0,2\pi]} \left\{ \sum_{\lambda \in \Lambda} \widehat{\omega}_p((2N)^{-1}(\xi + 2\pi\lambda)) \overline{\widehat{\tilde{\omega}}_p((2N)^{-1}(\xi + 2\pi\lambda))} \right\} \\
 &\quad \times h_q((2N)^{-1}\xi) \overline{\tilde{h}_q((2N)^{-1}\xi)} e^{2\pi i \lambda \xi} d\xi \\
 &= \int_{2N[0,2\pi]} h_q((2N)^{-1}\xi) \overline{\tilde{h}_q((2N)^{-1}\xi)} e^{2\pi i \lambda \xi} d\xi
 \end{aligned}$$

$$\begin{aligned}
&= \int_{[0,2\pi]} \sum_{\sigma=0}^{2N-1} h_p((2N)^{-1}(\xi + 2\pi\sigma)) \overline{\tilde{h}_q((2N)^{-1}(\xi + 2\pi\sigma))} e^{2\pi i\lambda\xi} d\xi \\
&= \int_{[0,2\pi]} e^{2\pi i\lambda\xi} d\xi = \delta_{0,\lambda} \quad \square
\end{aligned}$$

Theorem 3.3. Suppose $\{\omega_n(x) : n \geq 0\}$ and $\{\tilde{\omega}_n(x) : n \geq 0\}$ are nonuniform wavelet packets with respect to a pair of biorthogonal scaling functions $\omega_0(x)$ and $\tilde{\omega}_0(x)$, respectively. Then, for $p \geq 0$, we have

$$\langle \omega_{q_1+2Np}(\cdot), \tilde{\omega}_{q_2+2Np}(\cdot - \lambda) \rangle = \delta_{0,\lambda} \delta_{q_1,q_2}, \quad \lambda \in \Lambda, 0 \leq q_1, q_2 \leq 2N - 1. \quad (3.3)$$

Proof. By Lemma 2.5, we have

$$\begin{aligned}
\langle \omega_{q_1+2Np}(\cdot), \tilde{\omega}_{q_2+2Np}(\cdot - \lambda) \rangle &= \langle \hat{\omega}_{q_1+2Np}(\cdot), \hat{\tilde{\omega}}_{q_2+2Np}(\cdot - \lambda) \rangle \\
&= \int_{\mathbb{R}} \hat{\omega}_{q_1+2Np}(\xi) \overline{\hat{\tilde{\omega}}_{q_2+2Np}(x)} e^{2\pi i\lambda\xi} d\xi \\
&= \int_{\mathbb{R}} h_{q_1}((2N)^{-1}\xi) \hat{\omega}_p((2N)^{-1}\xi) \overline{\tilde{h}_{q_2}((2N)^{-1}\xi)} \overline{\hat{\tilde{\omega}}_p((2N)^{-1}\xi)} e^{2\pi i\lambda\xi} d\xi \\
&= 2N \sum_{\lambda \in \Lambda} \int_{([0,2\pi]+2\pi\lambda)} h_{q_1}(\xi) \hat{\omega}_p(\xi) \overline{\tilde{h}_{q_2}(\xi)} \overline{\hat{\tilde{\omega}}_p(\xi)} e^{2\pi i(2N)\lambda\xi} d\xi \\
&= 2N \int_{[0,2\pi]} \sum_{\lambda \in \Lambda} \hat{\omega}_p(\xi - \lambda) \overline{\hat{\tilde{\omega}}_p(\xi - \lambda)} h_{q_1}(\xi) \overline{\tilde{h}_{q_2}(\xi)} e^{2\pi i(2N)\lambda\xi} d\xi \\
&= \int_{2N[0,2\pi]} h_{q_1}((2N)^{-1}\xi) \overline{\tilde{h}_{q_2}((2N)^{-1}\xi)} e^{2\pi i\lambda\xi} d\xi \\
&= \int_{[0,2\pi]} \sum_{\sigma=0}^{2N-1} h_{q_1}((2N)^{-1}(\xi + 2\pi\sigma)) \overline{\tilde{h}_{q_2}((2N)^{-1}(\xi + 2\pi\sigma))} e^{2\pi i\lambda\xi} d\xi \\
&= \int_{[0,2\pi]} \delta_{q_1,q_2} e^{2\pi i\lambda\xi} d\xi = \delta_{0,\lambda} \delta_{q_1,q_2}. \quad \square
\end{aligned}$$

Theorem 3.4. If $\{\omega_n(x) : n \geq 0\}$ and $\{\tilde{\omega}_n(x) : n \geq 0\}$ are basic nonuniform wavelet packets with respect to a pair of biorthogonal scaling functions $\omega_0(x)$ and $\tilde{\omega}_0(x)$, respectively. Then, for $\ell, n \geq 0$, we have

$$\langle \omega_\ell(\cdot), \tilde{\omega}_n(\cdot - \lambda) \rangle = \delta_{\ell,n} \delta_{0,\lambda}, \quad \lambda \in \Lambda. \quad (3.4)$$

Proof. For $\ell = n$, the result (3.4) follows by Theorem 3.2. When $\ell \neq n$, and $0 \leq \ell, n \leq 2N - 1$, the result (3.4) can be established from Theorem 3.3. Assuming that ℓ is not equal to n , and atleast one of $\{\ell, n\}$ does not lies in $0, 1, \dots, 2N - 1$, then we can rewrite ℓ, n as $\ell = q_1 + 2Np_1$, $n = s_1 + 2Nr_1$, where $p_1, r_1 \geq 0$, $0 \leq q_1, s_1 \leq 2N - 1$.

Case 1. If $p_1 = r_1$, then $q_1 \neq s_1$. Therefore, Eq. (3.4) follows by virtue of (2.19), (2.20), Lemma 2.5 and (3.1), i.e.,

$$\begin{aligned}
 \langle \omega_\ell(\cdot), \tilde{\omega}_n(\cdot - \lambda) \rangle &= \langle \omega_{q_1+2Np_1}(\cdot), \tilde{\omega}_{s_1+2Nr_1}(\cdot - \lambda) \rangle \\
 &= \langle \hat{\omega}_{q_1+2Np_1}(\cdot), \widehat{\tilde{\omega}}_{s_1+2Nr_1}(\cdot - \lambda) \rangle \\
 &= \int_{\mathbb{R}} \hat{\omega}_{q_1+2Np_1}(\xi) \overline{\widehat{\tilde{\omega}}_{s_1+2Nr_1}(\xi)} e^{2\pi i \lambda \xi} d\xi \\
 &= \int_{\mathbb{R}} h_{q_1}((2N)^{-1}\xi) \hat{\omega}_{p_1}((2N)^{-1}\xi) \overline{\widehat{\tilde{\omega}}_{r_1}((2N)^{-1}\xi) \tilde{h}_{s_1}((2N)^{-1}\xi)} e^{2\pi i \lambda \xi} d\xi \\
 &= \sum_{\lambda \in \Lambda} \int_{2N([0,2\pi]+\lambda)} h_{q_1}((2N)^{-1}\xi) \hat{\omega}_{p_1}((2N)^{-1}\xi) \overline{\widehat{\tilde{\omega}}_{r_1}((2N)^{-1}\xi)} \\
 &\hspace{20em} \times \overline{\tilde{h}_{s_1}((2N)^{-1}\xi)} e^{2\pi i \lambda \xi} d\xi \\
 &= \int_{2N[0,2\pi]} \left\{ \sum_{\lambda \in \Lambda} \hat{\omega}_{p_1}((2N)^{-1}(\xi + \lambda)) \overline{\widehat{\tilde{\omega}}_{r_1}((2N)^{-1}(\xi + \lambda))} \right\} \\
 &\hspace{10em} \times h_{q_1}((2N)^{-1}\xi) \overline{\tilde{h}_{s_1}((2N)^{-1}\xi)} e^{2\pi i \lambda \xi} d\xi \\
 &= \int_{[0,2\pi]} \sum_{\sigma=0}^{2N-1} h_{q_1}((2N)^{-1}(\xi + 2\pi\sigma)) \overline{\tilde{h}_{s_1}((2N)^{-1}(\xi + 2\pi\sigma))} e^{2\pi i \lambda \xi} d\xi \\
 &= \int_{[0,2\pi]} \delta_{q_1, s_1} e^{2\pi i \lambda \xi} d\xi = \delta_{0, \lambda} = 0.
 \end{aligned}$$

Case 2. If $p_1 \neq r_1$, order $p_1 = q_2 + 2N p_2$, $r_1 = s_2 + 2N r_2$, where $p_2, r_2 \geq 0$ and $0 \leq q_2, s_2 \leq 2N - 1$. If $p_2 = r_2$, then $q_2 \neq s_2$. Similar to Case 1, (3.4) can be established. When $p_2 \neq r_2$, we order $p_2 = q_3 + 2N p_3$, $r_2 = s_3 + 2N r_3$, where $p_3, r_3 \geq 0$ and $0 \leq q_3, s_3 \leq 2N - 1$. Thus, after taking finite steps (denoted by κ), we obtain $0 \leq p_\kappa, r_\kappa \leq 2N - 1$ and $0 \leq q_\kappa, s_\kappa \leq 2N - 1$. If $p_\kappa = r_\kappa$, then $q_\kappa \neq s_\kappa$. Similar to the Case 1, (3.4) follows. If $p_\kappa \neq r_\kappa$, then it gets from (2.9)–(2.12) that

$$\langle \omega_{p_\kappa}(\cdot), \tilde{\omega}_{r_\kappa}(\cdot - \lambda) \rangle = 0, \lambda \in \Lambda \iff \sum_{\lambda \in \Lambda} \hat{\omega}_{p_\kappa}(\xi - \lambda) \overline{\widehat{\tilde{\omega}}_{r_\kappa}(\xi - \lambda)} = 0, \xi \in \mathbb{R}.$$

Furthermore, we obtain

$$\begin{aligned}
 \langle \omega_p(\cdot), \tilde{\omega}_r(\cdot - \lambda) \rangle &= \langle \hat{\omega}_p(\cdot), \widehat{\tilde{\omega}}_r(\cdot - \lambda) \rangle \\
 &= \langle \hat{\omega}_{q_1+2Np_1}(\cdot), \widehat{\tilde{\omega}}_{s_1+2Nr_1}(\cdot - \lambda) \rangle \\
 &= \int_{\mathbb{R}} \hat{\omega}_{q_1+2Np_1}(\xi) \overline{\widehat{\tilde{\omega}}_{s_1+2Nr_1}(x)} e^{2\pi i \lambda \xi} d\xi
 \end{aligned}$$

$$\begin{aligned}
&= \int_{\mathbb{R}} h_{q_1}((2N)^{-1}\xi) h_{q_2}((2N)^{-2}\xi) \widehat{\omega}_{p_2}((2N)^{-2}\xi) \overline{\widehat{\omega}_{r_2}((2N)^{-2}\xi)} \\
&\quad \times \overline{\widetilde{h}_{s_1}((2N)^{-1}\xi)} \overline{\widetilde{h}_{s_2}((2N)^{-2}\xi)} e^{2\pi i \lambda \xi} d\xi \\
&= \int_{\mathbb{R}} \left\{ \prod_{\ell=1}^{\kappa} h_{q_\ell}((2N)^{-\ell}\xi) \right\} \widehat{\omega}_{p_\kappa}((2N)^{-\kappa}\xi) \overline{\widehat{\omega}_{r_\kappa}((2N)^{-\kappa}\xi)} \left\{ \prod_{\ell=1}^{\kappa} \overline{\widetilde{h}_{s_\ell}((2N)^{-\ell}\xi)} \right\} e^{2\pi i \lambda \xi} d\xi \\
&= \sum_{\lambda \in \Lambda} \int_{(2N)^\kappa([0, 2\pi] + \lambda)} \left\{ \prod_{\ell=1}^{\kappa} h_{q_\ell}((2N)^{-\ell}\xi) \right\} \left\{ \widehat{\omega}_{p_\kappa}((2N)^{-\kappa}\xi) \overline{\widehat{\omega}_{r_\kappa}((2N)^{-\kappa}\xi)} \right\} \\
&\quad \times \left\{ \prod_{\ell=1}^{\kappa} \overline{\widetilde{h}_{s_\ell}((2N)^{-\ell}\xi)} \right\} e^{2\pi i \lambda \xi} d\xi \\
&= \int_{(2N)^\kappa[0, 2\pi]} \left\{ \prod_{\ell=1}^{\kappa} h_{q_\ell}((2N)^{-\ell}\xi) \right\} \left\{ \prod_{\ell=1}^{\kappa} \overline{\widetilde{h}_{s_\ell}((2N)^{-\ell}\xi)} \right\} \\
&\quad \times \left\{ \sum_{\lambda \in \Lambda} \widehat{\omega}_{p_\kappa}((2N)^{-\kappa}(\xi + \lambda)) \overline{\widehat{\omega}_{r_\kappa}((2N)^{-\kappa}(\xi + \lambda))} \right\} e^{2\pi i \lambda \xi} d\xi \\
&= \int_{(2N)^\kappa[0, 2\pi]} \left\{ \prod_{\ell=1}^{\kappa} h_{q_\ell}((2N)^{-\ell}\xi) \right\} \cdot 0 \cdot \left\{ \prod_{\ell=1}^{\kappa} \overline{\widetilde{h}_{s_\ell}((2N)^{-\ell}\xi)} \right\} e^{2\pi i \lambda \xi} d\xi \quad \square
\end{aligned}$$

4. The nonuniform biorthogonal wavelet packet bases of $L^2(\mathbb{R})$

In this section, we will decompose subspaces V_j, \widetilde{V}_j and W_j, \widetilde{W}_j by constructing a series of subspaces of nonuniform wavelet packets. Furthermore, we present the direct decomposition for space $L^2(\mathbb{R})$.

For any $n \geq 0$, define

$$U_n = \overline{\text{span}} \{ \omega_n(x - \lambda) : \lambda \in \Lambda \}, \quad (4.1)$$

Then, we have $U_0 = V_0$, $U_s = W_0^q$, $1 \leq q \leq 2N - 1$. Assume that $\{h_q((2N)^{-1}(\xi - \lambda))\}_{q, \lambda=0}^{2N-1}$ is a unitary matrix.

Lemma 4.1. For $n \geq 0$, the space $\mathcal{D}U_n$ can be decomposed into direct sum of U_{q+2Nn} , $0 \leq q \leq 2N - 1$, i.e.

$$\mathcal{D}U_n = \bigoplus_{q=0}^{2N-1} U_{q+2Nn}, \quad (4.3)$$

where \mathcal{D} is the dilation operator with respect to the dilation $2N$.

Proof. First, we claim that

$$\mathcal{D}U_n = \left\{ f(x) : f(x) = \sum_{q=0}^{2N-1} \sum_{\lambda \in \Lambda} b_\lambda^q \omega_{q+2Nn}(x - \lambda), \quad \sum_{\lambda \in \Lambda} |b_\lambda^q|^2 < \infty \right\}. \quad (4.4)$$

As for any $0 \leq q \leq 2N - 1$, by (2.17) and (4.1), $\omega_{q+2Nn}(x - \lambda) \in \mathcal{DU}_n$. Assume that $f(x) \in \mathcal{DU}_n$, then there exists a sequence $\{c_\lambda\}_{\lambda \in \Lambda}$ such that

$$f(x) = \sum_{\lambda \in \Lambda} c_\lambda \omega_n(2Nx - \lambda). \tag{4.5}$$

Further, if there exists a sequence $\{b_\lambda^q\}_{\lambda \in \Lambda}$, $0 \leq q \leq 2N - 1$, as for $f(x) \in \mathcal{DU}_n$, such that

$$f(x) = \sum_{q=0}^{2N-1} \sum_{\lambda \in \Lambda} b_\lambda^q \omega_{q+2Nn}(x - \lambda). \tag{4.6}$$

Taking Fourier transform on the both sides of (4.5) and (4.6), respectively and by using (2.19), we obtain

$$\hat{f}(\xi) = m((2N)^{-1}\xi) \hat{\omega}_n((2N)^{-1}\xi) = \sum_{q=0}^{2N-1} g_q(\xi) h_q((2N)^{-1}\xi) \hat{\omega}_n((2N)^{-1}\xi), \tag{4.7}$$

where

$$m(\xi) = \sum_{\lambda \in \Lambda} c_\lambda e^{-2\pi i \lambda \xi}, \quad g_q(\xi) = \sum_{\lambda \in \Lambda} b_\lambda^q e^{-2\pi i \lambda \xi}, \quad 0 \leq q \leq 2N - 1, \xi \in \mathbb{R}.$$

The above result (4.7) follows if the following equality holds:

$$m((2N)^{-1}\xi) = \sum_{q=0}^{2N-1} g_q(\xi) h_q((2N)^{-1}\xi). \tag{4.8}$$

For any sequence $\{c_\lambda\}_{\lambda \in \Lambda}$, we will prove that there exists a sequence $\{b_\lambda^q\}_{\lambda \in \Lambda}$, $0 \leq q \leq 2N - 1$

such that (4.8) is satisfied. Moreover, Eq. (4.8) is equivalent to the following equation:

$$m((2N)^{-1}(\xi - \lambda)) = \sum_{q=0}^{2N-1} g_q(\xi) h_q((2N)^{-1}(\xi - \lambda)), \quad 0 \leq \lambda \leq 2N - 1. \tag{4.9}$$

The solvability of Eq. (4.9) for every sequence $\{c_\lambda\}_{\lambda \in \Lambda}$ follows from the fact that the matrix $\{h_q((2N)^{-1}(\xi - \lambda))\}_{0 \leq \lambda, q \leq 2N-1}$ is a unitary matrix (see [6, 7]). Hence, equality (4.4) follows. Furthermore, applying Theorem 3.3, it follows that

$$\{ \omega_{q+2Nn}(x - \lambda) : n \geq 0, 0 \leq q \leq 2N - 1, \lambda \in \Lambda \},$$

forms a Riesz basis of \mathcal{DU}_n . □

Similar to (4.3), we can establish the following result:

$$\tilde{U}_0 = \tilde{V}_0, \quad \tilde{U}_s = \tilde{W}_0^q, \quad 0 \leq q \leq 2N - 1,$$

and

$$\mathcal{D}\tilde{U}_n = \bigoplus_{q=0}^{2N-1} \tilde{U}_{q+2Nn}. \tag{4.10}$$

For $\ell \in \mathbb{N}$, define $\tilde{\Gamma}_\ell = \sum_{j=0}^\ell (2N)^j \mu_j$, $\Gamma_\ell = \tilde{\Gamma}_\ell - \tilde{\Gamma}_{\ell-1}$. In what follows, we will give the direct decomposition of space $L^2(\mathbb{R})$.

Theorem 4.2. The family of functions $\{\omega_n(x - \lambda) : n \in \Gamma_\ell, \lambda \in \Lambda\}$ constitutes Riesz basis of $\mathcal{D}^\ell W_0$. In particular, $\{\omega_n(x - \lambda) : n \geq 0, \lambda \in \Lambda\}$ constitutes Riesz basis of $L^2(\mathbb{R})$.

Proof. By equation (4.3), we have $\mathcal{DU}_0 = \bigoplus_{q=0}^{2N-1} U_q$ i.e. $\mathcal{DU}_0 = U_0 \bigoplus_{q=1}^{2N-1} U_q$. Since $U_0 = V_0$ and $W_0 = \bigoplus_{q=1}^{2N-1} W_0^q = \bigoplus_{q=1}^{2N-1} U_q$, then $\mathcal{DU}_0 = V_0 \bigoplus W_0$. It can be inductively inferred from (4.3) that

$$\mathcal{D}^\ell U_0 = \mathcal{D}^{\ell-1} U_0 \bigoplus_{n \in \Gamma_\ell} U_n, \quad \ell \in \mathbb{N}. \tag{4.11}$$

Since $V_{j+1} = V_j \oplus W_j$, $j \in \mathbb{Z}$, therefore, $\mathcal{D}^\ell U_0 = \mathcal{D}^{\ell-1} U_0 \oplus \mathcal{D}^{\ell-1} W_0$, $\ell \in \mathbb{N}$. It follows from (4.3) and Proposition 2.4 that $\mathcal{D}^\ell W_0 = \bigoplus_{n \in \Gamma_\ell} U_n$ and

$$L^2(\mathbb{R}) = V_0 \oplus \left(\bigoplus_{\ell \geq 0} \mathcal{D}^\ell W_0 \right) = U_0 \oplus \left(\bigoplus_{\ell \geq 0} \left(\bigoplus_{n \in \Gamma_\ell} U_n \right) \right) = \bigoplus_{n=0}^{\infty} U_n. \quad (4.12)$$

In the light of Theorem 3.2, the family $\{\omega_n(x - \lambda) : \lambda \in \Lambda\}$ is a Riesz basis of $\mathcal{D}W_0$. Moreover, according to (4.12), the family $\{\omega_n(x - \lambda) : n \geq 0, \lambda \in \Lambda\}$ forms a Riesz basis of $L^2(\mathbb{R})$. \square

Corollary 4.3. For every $\ell \in \mathbb{N}$, the family of functions

$$\{\tilde{\omega}_n(x - \lambda) : n \in \Gamma_\ell, \lambda \in \Lambda\}$$

forms a Riesz basis of $\mathcal{D}^\ell \tilde{W}_0$.

Corollary 4.4. For every $\ell \in \mathbb{N}$, the family of functions

$$\{\omega_n((2N)^j x - \lambda) : j \in \mathbb{Z}, n \in \Gamma_\ell, \lambda \in \Lambda\}$$

forms a Riesz basis of $L^2(\mathbb{R})$.

5. References

- [1] B. Behera, Wavelet packets associated with nonuniform multiresolution analyses, *J. Math. Anal. Appl.* **328** (2007) 1237-1246.
- [2] Q. Chen and Z. Chang, A study on compactly supported orthogonal vector valued wavelets and wavelet packets, *Chaos, Solitons and Fractals*, **31** (2007) 1024-1034.
- [3] C. Chui and C. Li, Non-orthogonal wavelet packets, *SIAM J. Math. Anal.* **24**(3) (1993) 712-738.
- [4] A. Cohen and I. Daubechies, On the instability of arbitrary biorthogonal wavelet packets, *SIAM J. Math. Anal.* **24**(5) (1993) 1340-1354.
- [5] R. R. Coifman, Y. Meyer, S. Quake, and M. V. Wickerhauser, Signal processing and compression with wavelet packets, *Technical Report*, Yale University (1990).
- [6] J. P. Gabardo and M. Nashed, Nonuniform multiresolution analyses and spectral pairs, *J. Funct. Anal.* **158** (1998) 209-241.
- [7] J. P. Gabardo and M. Nashed, An analogue of Cohen's condition for nonuniform multiresolution analyses, in: A. Aldroubi, E. Lin (Eds.), *Wavelets, Multiwavelets and Their Applications*, in: *Cont. Math.*, **216**, Amer. Math. Soc. Providence, RI, (1998) 41-61.
- [8] J. P. Gabardo and X. Yu, Wavelets associated with nonuniform multiresolution analyses and one-dimensional spectral pairs, *J. Math. Anal. Appl.* **323** (2006) 798-817.
- [9] J. P. Gabardo and X. Yu, Nonuniform wavelets and wavelet sets related to one-dimensional spectral pairs, *J. Approx. Theory*, **145** (2007) 133-139.
- [10] J. Han and Z. Cheng, On the splitting trick and wavelets packets with arbitrary dilation matrix of $L^2(\mathbb{R}^s)$, *Chaos, Solitons and Fractals*, **40** (2009) 130-137.
- [11] S. G. Mallat, Multiresolution approximations and wavelet orthonormal bases of $L^2(\mathbb{R})$, *Trans. Amer. Math. Soc.* **315** (1989) 69-87.

- [12] F. A. Shah, Construction of wavelet packets on p -adic field, *Int. J. Wavelets Multiresolut. Inf. Process.* **7**(5) (2009) 553-565.
- [13] F. A. Shah and L. Debnath, p -Wavelet frame packets on a half-line using the Walsh-Fourier transform, *Integ. Transf. Special Funct.* **22**(12) (2011) 907-917.
- [14] Z. Shen, Non-tensor product wavelet packets in $L^2(\mathbb{R}^s)$, *SIAM J. Math. Anal.* **26**(4) (1995) 1061-1074.