

## Doubly Periodic Solution for Nonlinear Schrodinger's Equation with Triple Power Law Nonlinearity

Daniela Milovic<sup>1</sup>, Anjan Biswas<sup>2</sup> \*

<sup>1</sup>Faculty of Electronic Engineering, Department of Telecommunications, University of Nis, Aleksandra Medvedeva 14, 1800 Nis, SERBIA

<sup>2</sup> Center for Research and Education in Optical Sciences and Applications, Department of Applied Mathematics and Theoretical Physics, Delaware State University, Dover, DE 19901-2277 USA

(Received 20 May 2008, accepted 7 July 2008)

**Abstract:** The nonlinear Schrodinger's equation with triple-power law nonlinearity is studied in this paper. The travelling wave ansatz is used and the doubly periodic wave solution is obtained. The numerical simulation is also given at the end of this paper

**Keywords:** nonlinear Schrodinger's equation; elliptic functions; conservation laws; travelling waves

**AMS Codes:** 78A60, 37K10; 35Q51, 35Q55

**PACS Codes:** 02.30.Ik, 02.30.Jr, 42.81.Dp, 52.35.Sb

**OCIS Codes:** 060.2310; 060.4510; 060.5530; 190.3270; 190.4370

### 1 Introduction

The nonlinear Schrodinger's equation (NLSE) plays a vital role in various areas of physical, biological and engineering sciences. It appears in many applied fields like Fluid Dynamics, Nonlinear Optics, Plasma Physics and Protein Chemistry. The NLSE that is going to be studied in this book is given by [1-15].

$$iq_t + \frac{1}{2}q_{xx} + F(|q|^2)q = 0 \quad (1)$$

In (1),  $F$  is a real-valued algebraic function where  $F(|q|^2)q : C \mapsto C$ . Considering the complex plane  $C$  as a two-dimensional linear space  $R^2$ , it can be said that the function  $F(|q|^2)q$  is  $k$  times continuously differentiable so that one can write [10]

$$F(|q|^2)q \in \bigcup_{m,n=1}^{\infty} C^k((-n, n) \times (-m, m); R^2) \quad (2)$$

In Eq.(1),  $q$  is the dependent variable while  $x$  and  $t$  are the independent variables that represents space and time respectively. The first term in (1) represents the time evolution term while the second term is due to the group velocity dispersion and the third term accounts for nonlinearity. This is a nonlinear partial differential equation that is not integrable, in general. The nonintegrability is not necessarily related to the nonlinear term in (1). Higher order dispersion, for example, can also make the system non-integrable while it still remains Hamiltonian.

Eq.(1) governs the evolution of a wave packet in a weakly nonlinear and dispersive medium and has thus arisen in diverse fields such as water waves, plasma physics, nonlinear optics. One other application of

\*Corresponding author. E-mail address: biswas.anjan@gmail.com

this equation is in pattern formation, where it has been used to model some nonequilibrium pattern forming systems. In particular, this equation is now widely used in the optics field as a good model for optical pulse propagation in nonlinear fibers. It is also known to support soliton solutions for particular forms of the nonlinearity in  $F$ . In this paper, however, the interest is going to be on obtaining the doubly periodic solution to (1) by the travelling wave ansatz.

### 1.1 Integrals of motion

An important property of nonlinear evolution equations is that it has conserved quantities also known as *Integrals of Motion*. In fact, equation (1) has three integrals of motion. They are the energy ( $E$ ), linear momentum ( $M$ ) and Hamiltonian ( $H$ ) which are respectively given by

$$E = \int_{-\infty}^{\infty} |q|^2 dx \quad (3)$$

$$M = i \int_{-\infty}^{\infty} (q^* q_x - q q_x^*) dx \quad (4)$$

and

$$H = \int_{-\infty}^{\infty} \left[ \frac{1}{2} |q_x|^2 - f(I) \right] dx \quad (5)$$

where

$$f(I) = \int_0^I F(\xi) d\xi \quad (6)$$

with the intensity  $I = |q|^2$ . The first conserved quantity is also known as the mass, wave action or plasmon number; in optics, however, it is called the *wave power* while mathematically, it is known as the  $L_2$  norm. The Hamiltonian is one of the most fundamental notions in mechanics and more generally in the theory of conservative dynamical systems with finite or even infinite degrees of freedom. The most useful approach in the soliton theory of conservative non-integrable Hamiltonian system is a representation on the plane of conserved quantities namely the Hamiltonian-versus-energy diagrams [4]. One can see that (1) can now be written in a canonical form as

$$iq_t = \frac{\delta H}{\delta q^*} \quad (7)$$

$$iq_t^* = -\frac{\delta H}{\delta q} \quad (8)$$

where in (7) and (8) the right sides denote the Fréchet derivative

### 1.2 Travelling waves

A travelling wave is a solution of the NLSE that represents a wave of permanent form which does not change its shape during propagation and moves with a constant speed. The wave may be localized or periodic. In order to seek a travelling wave solution of the NLSE that is given by (1), introduce the ansatz

$$q(x, t) = Ag[B(x - \bar{x}(t))]e^{i(-\kappa x + \omega t + \sigma_0)} \quad (9)$$

where the function  $g$  represents the shape of the wave form described by (1) and it depends on the type of nonlinearity in it. Also, in (9)  $A$  and  $B$  respectively represent the amplitude and width of the wave,  $\kappa$  is the wave frequency while  $\omega$  is the wave number,  $\sigma_0$  is the phase while  $\bar{x}$  gives the mean position, so that the velocity of the soliton is given by

$$v = \frac{d\bar{x}}{dt} \quad (10)$$

The soliton width and the amplitude are related as  $B = \lambda(A)$  where the functional form  $\lambda$  depends on the type of nonlinearity in (1). Thus, from (9), one easily gets

$$q_t = -ABvg'(\tau)e^{i\phi} - i\omega Ag(\tau)e^{i\phi} \quad (11)$$

where, the phase  $\phi$  is

$$\phi(x, t) = -\kappa x + \omega t + \sigma_0 \quad (12)$$

and

$$\tau = B(x - \bar{x}) \quad (13)$$

Also, from (9) one can get

$$q_{xx} = AB^2g''(\tau)e^{i\phi} - 2i\kappa ABg'(\tau)e^{i\phi} - \kappa^2 Ag(\tau)e^{i\phi} \quad (14)$$

Substituting (11) and (14) into (1) simplifies it to

$$-iBvg' + \omega g + \frac{1}{2}B^2g'' - i\kappa Bg' - \frac{1}{2}\kappa^2 g + gF(A^2g^2) = 0 \quad (15)$$

From (15), equating the real and imaginary parts yields

$$ABg'(\kappa + v) = 0 \quad (16)$$

and

$$B^2g'' - (\kappa^2 - 2\omega)g + 2gF(A^2g^2) = 0 \quad (17)$$

so that from (16), one obtains

$$\kappa = -v \quad (18)$$

Now, multiplying both sides of (17) by  $g'$ , then integrating and choosing the integration constant to be zero since the wave profile is such that  $q$ ,  $q_x$  and  $q_{xx}$  approach zero as  $|x| \rightarrow \infty$  gives

$$B^2 (g')^2 - (\kappa^2 - 2\omega) g^2 + 2 \int (g^2)' F(A^2g^2) dg = 0 \quad (19)$$

On separation of variables and integrating once more, Eq.(19) leads to

$$x - vt = \int \frac{dg}{[(\kappa^2 - 2\omega)g^2 - 2 \int (g^2)' F(A^2g^2)]^{\frac{1}{2}}} \quad (20)$$

Eq.(20) can be integrated if the law of nonlinearity is known. This integral will be evaluated for the case of triple power law nonlinearity as will be seen in the following section.

## 2 Triple power law

In this paper, the type of nonlinearity that is going to be studied is given by triple-power law nonlinearity that is given by

$$F(s) = s^p + \nu_1 s^{2p} + \nu_2 s^{3p} \quad (21)$$

so that the NLSE that is being studied in this paper takes the form

$$iq_t + \frac{1}{2}q_{tt} + \left(|q|^{2p} + \nu_1 |q|^{4p} + \nu_2 |q|^{6p}\right)q = 0 \quad (22)$$

For this law of nonlinearity, although the energy and the linear momentum stays the same, the Hamiltonian takes the form

$$H = \int_{-\infty}^{\infty} \left( \frac{1}{2} |q_x|^2 - \frac{1}{p+1} |q|^{2p+2} - \frac{\nu_1}{2p+1} |q|^{4p+2} - \frac{\nu_2}{3p+1} |q|^{6p+2} \right) dx \tag{23}$$

It is to be noted that the case where  $\nu_2 = 0$ , with  $\nu_1 \neq 0$  is known as dual-power law nonlinearity. This case is studied in details in the context of nonlinear fiber optics [5].

For this kind of nonlinearity, (20) reduces to

$$x - vt = B \int \frac{\sqrt{(p+1)(2p+1)(3p+1)}}{g\sqrt{P}} dg \tag{24}$$

where

$$P = (\kappa^2 - 2\omega) (p+1)(2p+1)(3p+1) - 2A^{2p}g^{2p}(2p+1)(3p+1) - 2\nu_1A^{4p}g^{4p}(p+1)(3p+1) - 2\nu_2A^{6p}g^{6p}(p+1)(2p+1) \tag{25}$$

which integrates to

$$\begin{aligned} & (x - vt)p \sqrt{g^2 \left( \kappa^2 - 2\omega - 2A^{2p}g^{2p} - \frac{2\nu_1A^{4p}g^{4p}}{2p+1} - \frac{2\nu_2A^{6p}g^{6p}}{3p+1} \right) g_3} \\ &= \Pi \left[ 1 - \frac{g_2}{g_3}, \sin^{-1} \sqrt{\frac{g^{2p} - g_3}{g_2 - g_3}}, \frac{g_2 - g_3}{g_1 - g_3} \right] \sqrt{\frac{(g^{2p} - g_1)(g^{2p} - g_2)(g^{2p} - g_3)}{g_3 - g_1}} \end{aligned} \tag{26}$$

where  $\Pi$  is the incomplete elliptic integral of the third kind that is defined as

$$\Pi(n; \phi, \alpha) = \int_0^\phi \frac{d\theta}{\sqrt{1 - n \sin^2 \theta} \sqrt{1 - \sin^2 \alpha \sin^2 \theta}} \tag{27}$$

and

$$g_1 = \frac{1}{6\nu_2A^{6p}(2p+1)} \left[ -2\nu_1A^{4p}(3p+1) + \frac{2^{\frac{5}{3}}A^{8p}(3p+1) \{ \nu_1^2(3p+1) - 3\nu_2(2p+1)^2 \}}{R^{\frac{1}{3}}} + 2^{\frac{1}{3}}R^{\frac{1}{3}} \right] \tag{28}$$

$$g_2 = \frac{1}{12\nu_2A^{6p}(2p+1)} \left[ -4\nu_1A^{4p}(3p+1) + \frac{2^{\frac{5}{3}}(1 + i\sqrt{3})A^{8p}(3p+1) \{ 3\nu_2(2p+1)^2 - \nu_1^2(3p+1) \}}{R^{\frac{1}{3}}} + 2^{\frac{1}{3}}i(i + \sqrt{3})R^{\frac{1}{3}} \right] \tag{29}$$

$$g_3 = \frac{1}{12\nu_2A^{6p}(2p+1)} \left[ -4\nu_1A^{4p}(3p+1) + \frac{2^{\frac{5}{3}}i(i + \sqrt{3})A^{8p}(3p+1) \{ \nu_1^2(3p+1) - 3\nu_2(2p+1)^2 \}}{R^{\frac{1}{3}}} + 2^{\frac{1}{3}}(1 + i\sqrt{3})R^{\frac{1}{3}} \right] \tag{30}$$

with  $R$  being given by

$$\begin{aligned} R = & A^{12p} \left( -4\nu_1^3 + 18\nu_1\nu_2 + 27\kappa^2\nu_2^2 - 36p\nu_1^3 + 180p\nu_1\nu_2 + 243p\kappa^2\nu_2^3 \right. \\ & - 108p^2\nu_1^3 + 666p^2\nu_1\nu_2 + 810p^2\kappa^2\nu_2^2 - 108p^3\nu_1^3 + 1080p^3\nu_1\nu_2 + 1188p^3\kappa^2\nu_2^3 \\ & + 648p^4\kappa^2\nu_2^2 - 54\omega\nu_2^2 - 486p\omega\nu_2^2 - 1620p^2\omega\nu_2^2 - 2376p^3\omega\nu_2^2 - 1296p^4\omega\nu_2^2 \\ & \left. + (3p+1)A^{12p} \left[ 16(3p+1) \{ 3\nu_2(2p+1)^2 - \nu_1^2(3p+1) \}^3 \right. \right. \\ & \left. \left. + \{ 18\nu_1\nu_2(2p+1)^2(3p+1) - 4\nu_1^3(3p+1)^2 + 27\nu_2^2(2p+1)^3 (\kappa^2 - 2\omega) \}^2 \right]^{\frac{1}{2}} \right) \end{aligned} \tag{31}$$

This solution was plotted numerically and is shown in the following figure below. The solution represents the wave profile  $|q(x, t)|^2$  for  $p = 3/2$  and  $x$  is between  $-0.1$  and  $0.1$  while  $t$  varies from  $0.0$  to  $2.0$ .




Figure 1:  $|q(x, t)|^2$  for  $p = 3/2$  and  $x$  is between  $-0.1$  and  $0.1$  while  $t$  varies from  $0.0$  to  $2.0$ .

### 3 Conclusions

In this paper, the travelling wave solution to the NLSE with triple power law nonlinearity is studied. The doubly periodic wave solution is obtained and that result is plotted numerically. In future, this result will be extended to obtain soliton solutions and those results will be published elsewhere.

### Acknowledgement

The research work of the second author (AB) is funded by NSF-CREST Grant No: HRD-0630388 and the support is genuinely and sincerely appreciated.

### References

- [1] M. J. Ablowitz , H. Segur: Solitons and Inverse Scattering Transform. *SIAM. Philadelphia, PA, USA.* (1981)
- [2] N. N. Akhmediev ,A. Ankiewicz:Solitons Nonlinear Pulses and Beams. *Chapman and Hall, UK.* (1997)
- [3] N. N. Akhmediev:Spatial solitons in Kerr and Kerr-like media. *em Optical and Quantum Electronics.* 30(7-10):535-569(1998)
- [4] N. N. Akhmediev, A. Ankiewicz , R. Grimshaw: Hamiltonian-versus-energy diagrams in soliton theory. *Physical Review E.* 59(5):6088-6096(1999)
- [5] A. Biswas ,S. Konar: Introduction to non-Kerr law optical solitons. *CRC Press, Boca Raton, FL, USA.* (2006)
- [6] G. Fang, W. Chen ,Z. Hu:Well-posedness of the solution to D-P equation with dispersive term.*International Journal of Nonlinear Science.* 5(2):125-132(2008)
- [7] A. Hasegawa ,Y. Kodama:Solitons in Optical Communications.*Oxford University Press, UK.* (1995)

- [8] Y. S. Kivshar , B. A. Malomed: Dynamics of solitons in nearly integrable systems. *Rev. Mod. Phys.* 61: 763-915(1989)
- [9] L. Tian ,Y. Wang: Global conservative solutions of the generalized Camassa-Holm equation. *International Journal of Nonlinear Science.* 5(3): 195-202(2008)
- [10] S. Wabnitz, Y. Kodama ,A. B. Aceves : Control of Optical Soliton Interactions. *Optical Fiber Technology.* 1:(187-217)(1995)
- [11] L. Wang: New exact solutions for the MNV equation. *International Journal of Nonlinear Science.* 5(2):152-157(2008)
- [12] S. Wen, W. Zhang , C. Niu: On the well-posedness of the Degasperis-Procesi equation with the dispersive term. *International Journal of Nonlinear Science.* 5(2):170-177(2008)
- [13] S. Yu, W. Yang, J. Ruan, L. Tian: Solitary wave solutions to approximate fully nonlinear double sine-Gordon equation. *International Journal of Nonlinear Science.* 3(3):163-169(2007)
- [14] V. E. Zakharov ,E. A. Kuznetsov: Optical solitons and quasisolitons. *Journal of Experimental and Theoretical Physics.* 86(5):1035-1046(1998)
- [15] P. E. Zhidkov: *Korteweg-de Vries and Nonlinear Schrödinger Equations: Qualitative Theory* Springer Verlag. (2001)